

ADOLESCENT & YOUNG ADULT VAPING OVER TIME

A longitudinal survey of 752 adolescents/young adults in Manitoba

Julie-Anne McCarthy, Janique Fortier, Tamara Taillieu, Samantha Salmon, Ashley Stewart-Tufescu, Isabel Garcés Dávila, Harriet L. MacMillan, Jitender Sareen, Lil Tonmyr, Marni Brownell, Nathan C. Nickel, & Tracie O. Afifi



VAPING PREVALENCE

These findings show the need for and the importance of early prevention.



Scan here for full study



89% OF ADOLESCENTS WHO VAPED AT WAVE 1 CONTINUED TO VAPE ABOUT 1(+) YEAR LATER

These identified ACEs and risk factors can help inform programs, strategies, and potential groups to target for vaping interventions.

FACTORS ASSOCIATED WITH AN INCREASED LIKELIHOOD OF VAPING*

*Vaping at any time, Wave 1 only, Wave 2 only, and/or continuously



Foster care/
contact with
child protection



Parental
separation/
divorce



Adolescent &
household
mental illness



Emotional
abuse &
neglect



Parental
problems with
police



Exposure to
verbal intimate
partner violence



Household
substance use



Peer
victimization



Unsafe
neighbourhood



Parental
gambling



Parental
smoking/vaping